arXiv:hep-ph/9402244v1 7 Feb 1994

Feb 1994
McGill/94-11

A parity invariant $SU(3)_c \otimes SU(3)_L \otimes U(1)$ model

R. Foot[1]
Department of Physics, McGill University,
3600 University Street, Montreal, Quebec, Canada H3A 2T8.

ABSTRACT

We examine a $SU(3)_c \otimes SU(3)_L \otimes U(1)$ gauge model which has a parity symmetric Lagrangian. The parity symmetry has the novel feature that it interchanges the gluons with the $SU(3)_L$ gauge bosons (which contain the ordinary $SU(2)_L$ weak gauge bosons). We show that the model reduces to the standard model at low energies and also predicts new physics in the form of exotic fermions.

---

[1] email: Foot@hep.physics.mcgill.ca

The standard model has been rather successful in recent times. With the present lack of experimental guidence it is difficult to know what lies beyond the standard model. One possibility is that nature respects some type of symmetry not present in the standard model. In particular, discrete space-time symmetries (parity, time reversal and charge conjugation) may be symmetries of the Lagrangian describing nature, but which are not manifested in the standard model Lagrangian because it may be incomplete. In this sense, discrete space-time symmetries may have a *fundamental* role to play in nature even though they are not symmetries of the Standard model Lagrangian. It is an interesting possibility that nature is described by a Lagrangian which is invariant under one or all of the discrete space-time symmetries. One well known example is the usual left-right symmetric model with gauge group $SU(3)_c \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)$ and is invariant under a parity symmetry [1]. This parity symmetry interchanges the $SU(2)_L$ gauge bosons with the $SU(2)_R$ gauge bosons and is assumed to be spontaneously broken by the vacuum. The usual left-right symmetric model is not the only possible model which allows parity to be a symmetry of the Lagrangian. For example, the quark-lepton symmetry model has gauge group $SU(3)_q \otimes SU(3)_l \otimes SU(2)_L \otimes U(1)$ and has a discrete symmetry which interchanges quarks and leptons [2]. This discrete symmetry can also be interpreted as parity (which is assumed to be spontaneously broken by the vacuum) and is thus another realization of a gauge model with left-right symmetry spontaneously broken [3].

The purpose of this letter is to point out that there is another type of model with left-right symmetry which is spontaneously broken. The idea is to build a model in which parity interchanges the gluons with the weak gauge bosons (W_L's) (rather than $W'_L s$ interchanging with $W'_R s$ as is the case in the usual left-right symmetric model). In-order for parity to interchange gluons with W_L's we need to enlarge the standard model gauge symmetry to:

$$SU(3)_c \otimes SU(3)_L \otimes U(1)_{y'} \tag{1}$$

We then need to construct the fermions so that they are symmetric when the two $SU(3)$'s are interchanged. The model must also reduce to the standard model at some scale if it is to describe the real world. This can be achieved with the following fermion representations, (under the gauge symmetry eq.(1)):

$$\begin{array}{lll} f^a_L \sim (1, 3, -1/3), & f'^a_L \sim (1, 3, -1/3), & F^a_L \sim (1, 3, 2/3), \\ d^a_R \sim (3, 1, -1/3), & d'^a_R \sim (3, 1, -1/3), & u^a_R \sim (3, 1, 2/3), \\ Q^a_L \sim (3, \bar{3}, 0), & & \end{array} \tag{2}$$

where $a = 1, 2, 3$, is the generation index [4]. It is straightforward to check that with the above choice of gauge quantum numbers, all of the gauge anomalies cancel per generation. We now introduce the discrete parity operation which we define below:

$$x \to -x, \ t \to t,$$

$$G^a_{c\mu} \leftrightarrow W_L^{a\mu},\ B_\mu \leftrightarrow B^\mu,$$

$$f_L \leftrightarrow \gamma_0 d_R,\ f'_L \leftrightarrow \gamma_0 d'_R,\ u_R \leftrightarrow \gamma_0 F_L,\ Q_L \leftrightarrow \gamma_0 (Q_L)^c, \tag{3}$$

where $G^a_{c\mu}$, $W^a_{L\mu}$, B_μ are the gauge bosons of $SU(3)_c$, $SU(3)_L$ and $U(1)_{y'}$ respectively. It is straightforward to check that the gauge covariant kinetic terms of the fermions and the gauge bosons are invariant with respect to the parity symmetry (eq.(3)) provided that the bare gauge coupling constants of $SU(3)_c$ and $SU(3)_L$ are equal.

Gauge symmetry breaking to the standard model can be achieved by introducing a Higgs field, $\chi_1 \sim (1, \bar{3}, 1/3)$. The field χ_1 couples to the fermions of the model as follows:

$$\mathcal{L}^{\chi_1} = \lambda_1 \bar{Q}_L \chi_1 d'_R + \lambda_2 \bar{f}'_L \chi_1 (F_L)^c + H.c., \tag{4}$$

where we have suppressed the generation indicies. Provided that χ_1 develops the vacuum expectation value (VEV):

$$\langle \chi_1 \rangle = \left(\begin{array}{c} 0 \\ 0 \\ w \end{array} \right), \tag{5}$$

then the gauge symmetry breaks down to the Standard model:

$$\begin{array}{c} SU(3)_c \otimes SU(3)_L \otimes U(1)_{y'} \\ \downarrow \langle \chi_1 \rangle \\ SU(3)_c \otimes SU(2)_L \otimes U(1)_Y \end{array} \tag{6}$$

where $Y = 2y' - \lambda_8/\sqrt{3}$ is the combination of y' and λ_8 which is left unbroken by $\langle \chi_1 \rangle$ ($\lambda_8 = diag(1, 1, -2)/\sqrt{3}$). From eq.(2) it is easy to see that each generation contains 27 Weyl fermions in this model. One can identify 15 of these Weyl fermions as the standard quarks and leptons with the correct gauge quantum numbers under the $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ subgroup left unbroken by the VEV $\langle \chi_1 \rangle$. In particular, note that $Y(= 2y' - \lambda_8/\sqrt{3})$, is numerically identical to the standard model hypercharge assignments. In addition to these standard 15 Weyl fermions, there are 12 exotic fermion fields per generation. Under the $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ subgroup, these 12 exotic fermions have the following gauge quantum numbers: $d'_{L,R} \sim (3, 1, -2/3)$, $X'_{L,R} = (V_{L,R}, E_{L,R})^T \sim (1, 2, -1)$, $\nu_R, \nu'_R \sim (1, 1, 0)$. We identify an exotic $SU(2)_L$ singlet charged -1/3 quark, and a leptonic (i.e. color singlet) vector-like $SU(2)_L$ doublet of leptons, together with two singlet Weyl neutrinos. Observe that the exotic charged $-1/3$ quarks gain masses and the exotic lepton doublet gain masses from the Yukawa Lagrangian terms in eq.(4) when χ_1 develops the VEV $\langle \chi_1 \rangle$ (eq.(5)). In order to fascillitate understanding we can write the $SU(3)_L$ degree's of freedom explicitly:

$$f_L = \begin{pmatrix} \nu_L \\ e_L \\ (\nu_R)^c \end{pmatrix}, \quad f'_L = \begin{pmatrix} V_L \\ E_L \\ (\nu'_R)^c \end{pmatrix}, \quad F_L = \begin{pmatrix} (E_R)^c \\ (V_R)^c \\ (e_R)^c \end{pmatrix},$$
$$Q_L = \begin{pmatrix} d_L \\ u_L \\ d'_L \end{pmatrix}, \qquad d_R,\ u_R,\ d'_R \tag{7}$$

In order to preserve the parity symmetry in the Yukawa sector, we need to introduce a scalar mulitplet χ_2 in addition to χ_1. The scalar χ_2 has the "mirror" gauge quantum numbers i.e. $\chi_2 \sim (\bar{3}, 1, 1/3)$ and couples to the fermions as follows:

$$\mathcal{L}^{\chi_2} = \lambda_1 \overline{(Q_L)^c} \chi_2 f'_L + \lambda_2 \bar{d}'_R \chi_2 (u_R)^c + H.c. \tag{8}$$

Note that under parity, $\mathcal{L}^{\chi_1} \leftrightarrow \mathcal{L}^{\chi_2}$, and the total Lagrangian remains invariant. Also note that we must require χ_2 to have zero VEV if we want to keep $SU(3)_c$ unbroken.

Ordinary electroweak symmetry breaking and fermion mass generation can be achieved by introducing two $SU(3)_L$ triplets $\rho_1,\ \eta_1$:

$$\begin{aligned} \mathcal{L}^{\eta_1,\rho_1} = \ & \lambda'_1 \bar{Q}_L d_R \rho_1 + \lambda'_2 \bar{Q}_L d'_R \rho_1 + \lambda'_3 \bar{Q}_L u_R \eta_1 + \lambda'_4 \bar{f}_L (F_L)^c \rho_1 + \lambda'_5 \bar{f}'_L (F_L)^c \rho_1 \\ & + \lambda'_6 \bar{f}_L \eta_1 (f_L)^c + \lambda'_7 \bar{f}_L \eta_1 (f'_L)^c + + \lambda'_8 \bar{f}'_L \eta_1 (f'_L)^c + H.c. \end{aligned} \tag{9}$$

where the generation indices have been suppressed and $\rho_1 \sim (1, \bar{3}, 1/3),\ \eta_1 \sim (1, \bar{3}, -2/3)$. Note that we require that ρ_1, η_1 develop the VEVs:

$$\langle \rho_1 \rangle = \begin{pmatrix} u \\ 0 \\ 0 \end{pmatrix}, \ \langle \eta_1 \rangle = \begin{pmatrix} 0 \\ v \\ 0 \end{pmatrix}. \tag{10a, b}$$

These VEVs also give masses to the usual W, Z bosons, and the standard model intermediate gauge symmetry is broken in the usual way to $SU(3)_c \otimes U(1)_Q$:

$$\begin{array}{c} SU(3)_c \otimes SU(3)_L \otimes U(1)_{y'} \\ \downarrow \langle \chi_1 \rangle \\ SU(3)_c \otimes SU(2)_L \otimes U(1)_Y \\ \downarrow \langle \rho_1 \rangle, \langle \eta_1 \rangle \\ SU(3)_c \otimes U(1)_Q \end{array} \tag{11}$$

where $Q = I_3 + Y/2$, and is of course just the usual electric charge operator.

In order to maintain the discrete symmetry, the mirror partners ρ_2, η_2 also need to be introduced and as in the case of χ_2, they must gain zero VEV. It is easy to deduce their coupling to fermions, which is of course dictated by the requirement that the discrete symmetry be a symmetry of the Lagrangian:

$$\begin{aligned} \mathcal{L}^{\eta_2,\rho_2} = \ & \lambda'_1 \overline{(Q_L)^c} f_L \rho_2 + \lambda'_2 \overline{(Q_L)^c} f'_L \rho_2 + \lambda'_3 \overline{(Q_L)^c} F_L \eta_2 + \lambda'_4 \bar{d}_R (u_R)^c \rho_2 \\ & + \lambda'_5 \bar{d}'_R (u_R)^c \rho_2 + \lambda'_6 \bar{d}_R \eta_2 (d_R)^c + \lambda'_7 \bar{d}_R \eta_2 (d'_R)^c + + \lambda'_8 \bar{d}'_R \eta_2 (d'_R)^c + H.c. \end{aligned} \tag{12}$$

The existence of the discrete symmetry means that the bare $SU(3)_c$ and $SU(3)_L$ coupling constants must be equal to each other. This unfortunately tends to imply that $SU(3)_L$ symmetry breaking must occur at a very high energy scale (near 10^{15} GeV). This means that with the simplest symmetry breaking scenario given here, it won't be possible to detect any of the new gauge bosons predicted by the model. However, it may be possible to detect the exotic fermions or the effects of the mixing. The exotic charged $-1/3$ quarks can mix with the ordinary d quark, which will imply a small loss of GIM cancellation (since the exotic d' quark has non-canonical $SU(2)_L \otimes U(1)_Y$ gauge quantum numbers). The phenomenology of this has already been studied in the liturature [5].

The model can also incorporate neutrino masses. This may be important in the light of the solar neutrino problem [6]. One interesting point is that if $\lambda_6^{'} \gg \lambda_7^{'}$ (in eq.(9)), then two neutrino masses are nearly degenerate (in the limit that $\lambda_7^{'} = 0$, two of the ordinary weakly interacting neutrinos gain degerate Dirac masses while one remains massless [7]. In the MSW solution of the solar neutrino problem, the neutrino mass difference needs to be very small ($\sim 10^{-4}$ eV). This can be achieved naturally if either one neutrino is much heavier than the other or they are nearly degenerate. The latter case is perhaps phenomenologically more interesting because it means that the neutrino masses can be a lot larger than their mass difference and thus it may be possible to detect the neutrino masses directly in the Laboratory.

Finally note that the model has electric charge quantization classically [8]. By this, I mean that the classical structure of the Lagrangian such as the gauge invariance of the Yukawa Lagrangians (eq.(4) and eq.(9)), and the discrete parity symmetry imply that the $U(1)_{y'}$ charges of the fermions *and* the scalars are uniquely fixed (upto an overall normalization factor, which can be absorbed into the definition of the coupling constant). This fact then implies that the electric charges are also uniquely determined (up to an overall normalization factor), and thus every electric charge ratio is fixed in this theory.

In conclusion, we have proposed a simple gauge model which is defined by a Lagrangian with a novel realization of parity symmetry. The parity operation interchanges the gluons of QCD with the weak gauge bosons of an extended $SU(3)_L$ weak interaction. We have pointed out that the gauge model reduces successfully to the standard model at low energies. The principle phenomenological feature of the model is the prediction of new fermions. In particular exotic $SU(2)_L$ singlet charged -1/3 quarks and exotic leptons together with two right-handed neutrinos per generation are predicted. Neutrinos are expected to be massive in the model, and the model may be relevant to the MSW solution of the solar neutrino problem.

References

[1] J. C. Pati and A. Salam, Phys. Rev. D10, 1502 (1974); R. N. Mohapatra and J. C. Pati, Phys. Rev. D11, 566, 2558 (1974); G. Senjanovic and R. N. Mohapatra, Phys. Rev. D12, 1502 (1974).

[2] R. Foot and H. Lew, Phys. Rev. D41, 3502 (1990); R. Foot, H. Lew and R. R. Volkas, Mod. Phys. Lett. A8, 1859 (1993).

[3] It is also possible to define a gauge model in which parity is a symmetry of the Lagrangian and the symmetry is *not* broken by the vacuum but remains exact after symmetry breaking. See R. Foot, H. Lew and R. R. Volkas, Phys. Lett. B272, 67 (1991); Mod. Phys. Lett. A7, 2567 (1992).

[4] The model is related to the $SU(3)^3$ model (See e.g. A. de Rujula, H. Georgi and S. L. Glashow, in Fifth Workshop on Grand Unification, edited by K. Kang, H. Fried and P. Frampton, (World Scientific, Singapore, 1984), however that model does not have a Z_2 parity symmetric Lagrangian (although it does have a Z_3 symmetry). Some of the phenomenology of the two models is similiar, e.g. they have the same number of exotic fermions with the same gauge quantum numbers under the $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ subgroup. My model (like the $SU(3)^3$ model) is also related to E_6 models, and the 27 Weyl fermions in eq.(2) can fit into the 27 irreducible representation of E_6, however, again the E_6 model is explicitly parity violating. My motivation for the $SU(3)^2 \otimes U(1)$ model is that it is a new type of model which has a parity conserving Lagrangian.

[5] See for example, P. Langacker and D. London, Phys. Rev. D38, 866 (1988).

[6] For a review of the solar neutrino problem, see for example J. N. Bahcall, Neutrino astrophysics (CUP), 1989.

[7] To see this note that the λ_6' term gives rise to a 3×3 mass matrix which is antisymmetric in generation space. (This becomes evident by noting that under interchange of the two identical $f_L's$, the Lorentz contraction is antisymmetric, there is an antisymmetric factor because of Fermi-Dirac statistics, and there is a antisymmetry because of the $SU(3)_L$ contraction is antisymmetric. Thus only the anti-symmetric piece in generation space is non-vannishing.) A 3×3 antisymmetric mass matrix has two degenerate eigenvalues and one zero eigenvalue.

[8] For a review of electric charge quantization in the standard model and beyond, see R. Foot, G. C. Joshi, H. Lew and R. R. Volkas, Mod. Phys. Lett. A5, 2721 (1990).